



OVINE SEMEN SEXING AGENT- MALE

PRODUCT: RAMPLUS-20

PRODUCT #: RP-20

INDICATIONS

For increasing the percentage of MALE lambs in ovine species.

DOSAGE

- Each vial contains 20 units of sexing agent and will treat a total of 5 ml of extended semen. One unit will sex 0.25 ml of semen.
- Recommended insemination dose: 0.25 ml or 0.5 ml
- Recommended dilution: 5 ml of extended semen
- Recommended extension titer: 100-200 million live sperm/ml
- Recommended insemination titer : 50 million live sperm/ AI dose

NOTICE

RAMPLUS is a non-prescription biopharmaceutical agent. Federal law (US) does not require that this product be used by or on the order of a licensed veterinarian. This is not a restricted drug. Please consult local regulations.

DESCRIPTION

RAMPLUS is spermagenic agent for sexing of ovine semen. Packaged in kit form, each vial is lyophilized and sealed to maintain potency during storage. The agent is activated by the addition of extender directly to the RAMPLUS vial. After a short incubation in the RAMPLUS, the sexed semen is ready for cervical or laparoscopic artificial insemination (LAI).

MODE OF ACTION

RAMPLUS works by enhancing the fertility of the Y-chromosome bearing (male) sperm and reducing the fertility of the X-chromosome bearing (female) sperm. When inseminated near the time of ovulation, more ova are fertilized by the Y-chromosome bearing (male) sperm. The MALE SEX RATIO is increased to 70-80% and conception is boosted 5-25%.

FRESH SEMEN - PREPARATION

- After collection of semen, determine semen parameters.
- Dilute the appropriate volume of raw semen using a standard OVINE semen extender. See recommended extension rates.
- Equilibrate both the extended semen and the RAMPLUS-20 vial to 38°C (100.4°F) for 5-10 minutes.
- Add 5.0 ml of warm extended semen directly to the RAMPLUS-20 vial. Reseal the vial and gently mix.
- **IMPORTANT:** Incubate for **30 minutes** at 37-38°C (99-100°F).
- Load straws as usual and use immediately. See Table 1.
- Sexed semen can be cooled and stored according to standard procedures. Use within 24 hrs.

FROZEN SEMEN – PREPARATION

- Prepare and warm semen extender (7% glycerol) to 38°C (100.4°F). One-step TRIS-CITRATE based extenders are recommended.
- Extend semen by slow addition of warm extender to 80% final volume (4 ml). Equilibrate to 38°C (100.4°F). See recommended extension rates.
- Re-hydrate the RAMPLUS-20 vial with 1.0 ml of warm extender (100.4°F, 38°C). See Table 1.
- Add 4.0 ml. of extended semen to the warm RAMPLUS-20 vial. This will bring the extension to 100% final volume. Gently mix. See Table 1.
- **IMPORTANT:** Incubate for **30 minutes** at 38°C (100.4°F).
- Cool sexed semen to 5°C (42°F) over a period of 2.5 hours and allow to equilibrate for at least 4 hours prior to freezing.
- Load and seal straws as usual. Freeze following standard procedure.

Table 1 - Preparation of 5 ml of sexed semen*

Straw volume (ml)	No. of straws	RAMPLUS-20	Extended Semen	RAMPLUS SEMEN
0.25	20	1.0 ml	4.0 ml	5.0 ml
0.50	10	1.0 ml	4.0 ml	5.0 ml

- *Final concentration of 50 x 10⁶ live sperm/straw (pre-freeze)
- **Dilution ratio = 1 unit of RAMPLUS: 0.25 ml of extended semen

RECOMMENDED SEMEN EXTENSION

- **0.25 ml straws:** 20 straws x 50 million live sperm/straw (pre-freeze) / 20 straws x 0.25 ml/straw] = 1000 million live sperm/5 ml of extended semen = 200 million live sperm/ml.
- **0.50 ml straws:** 10 straws x 50 million live sperm/straw (pre-freeze) / 10 straws x 0.50 ml/straw] = 500 million live sperm/5 ml of extended semen = 100 million live sperm/ml.


TIMING OF INSEMINATION

Variations in synchronization protocols, superovulation regimens and breed differences will give rise to variations in the time of onset of heat and time of ovulation. These variances are important when using fixed-time breeding. For fixed-timed inseminations, please consult the following recommendations. In general, apply RAMPLUS 4-6 hours prior to ovulation. Ovulation occurs 24-27 hours after onset of heat or 65 hours after sponge removal.

1. Synchronized with progestin implant (CIDR®) + PMSG (PG 600®, Folligon®) or FSH: Fixed-timed inseminations: Breed 56 hours after progestin withdrawal. See Fig. 1A

2. Synchronized with progestin implant (pessary sponge) + PMSG (PG 600®, Folligon®) or FSH: Fixed-timed inseminations: Breed 66 hours after progestin withdrawal. See Fig. 1B.
3. Synchronized heats and natural heats: Estrus detection: Breed 32 hours after marking by vasectomized male.

Fig. 1. Fixed-timed estrus synchronization protocols for sheep.



STORAGE CONDITIONS

Store product in the freezer (-15-20°C). Keep desiccated and in the dark. Avoid moisture and strong sunlight. Do not reuse or store after dilution.

HOW SUPPLIED

RAMPLUS is lyophilized in the following package sizes: 0.25 ml, 0.5 ml, 10 and 20 unit vials.

WARNINGS

KEEP OUT OF REACH OF CHILDREN

Manufactured by:



EMLAB GENETICS, LLC
Arcola, IL 61910 USA

For more information, please call 708-442-3964 or log on to www.emlabgenetics.com

RAMPLUS™ is a trademark of **EMLAB GENETICS, LLC**

EMLAB GENETICS © 2018

REV 05.16.18